

Supplementary Information

Extraction and estimation of antistatic agent glycerol monostearate in polypropylene by gas chromatography coupled with flame ionisation detector

Sivakumar Padmanaban*, Vijay K Marrapu*, Rabindra Sahoo, Siva Jyothi P, and Hemant Tyagi

Quality Control Laboratory, Paradip Refinery, Indian Oil Corporation Limited, Paradip Refinery,
Paradip 754 141, Odisha, India

*E-mail: sivakumarp2@indianoil.in (SP), marrapuvk@indianoil.in (VKM)

Received 4 March 2022; accepted (revised) 2 January 2023

S. No	Contents	Pg. No.
1	Fig. S1 — GC-FID Chromatogram of GMS Standard 50 mg/L	2
2	Fig. S2 — GC-FID Chromatogram of GMS Standard 101.0 mg/L	3
3	Fig. S3 — GC-FID Chromatogram of GMS Standard 151.5 mg/L	4
4	Fig. S4 — Calibration overlay of lower concentrations 1.0, 2.5, 5.0 & 10.0 mg/L	5
5	Fig. S5 — Calibration overlay of lower concentrations 50.5, 101.0, 151.5 mg/L	6




Fig. S1 — GC-FID Chromatogram of GMS Standard 50 mg/L




Fig. S2 — GC-FID Chromatogram of GMS Standard 101.0 mg/L




Fig. S3 — GC-FID Chromatogram of GMS Standard 151.5 mg/L




Fig. S4 — Calibration overlay of lower concentrations 1.0, 2.5, 5.0 & 10.0 mg/L




Fig. S5 — Calibration overlay of lower concentrations 50.5, 101.0, 151.5 mg/L