

SHORT COMMUNICATION

A new antibacterial imidazole from the marine sponge *Ircinia fusca*

Srinu Meesala^{1*}, Ruby Singh¹, Milind G Watve¹ and Dattatraya G. Naik²

¹Department of Biology, Indian Institute of Science Education and Research, Dr. Homi Bhabha Road, Pune-411008, India

²Abasaheb Garware College, Karve Road, Pune, Maharashtra 411004, India

Received 29 May 2017; Revised 25 September 2017

A new imidazole alkaloid (1) along with two known compounds, variabilin (2) and iricinialactam A (3) have been isolated from the Arabian marine sponge *Ircinia fusca*. The structure of the new compound was established as 4-((1, 2-dihydroxy-5-(methyl (1-methyl-1H-imidazol-4-yl) amino) pentan-3-yl) oxy)-3, 5-dimethoxy-1-methylpyrrolidin-2-one (1) by 1D and 2D NMR, and high-resolution electrospray ionization mass spectrometry (HRESIMS). Compound 1 exhibited selective growth inhibitory activity against gram-positive bacteria *S. aureus* at 100 µg/mL.

Keywords: Anticancer activity, Antimicrobial, Imidazole alkaloids, Mass spectrometry, NMR, Sponge.

IPC code; Int. cl. (2015.01)–A01N 63/00, C07D 403/14

Introduction

Marine sponges are a rich source of metabolites reported in marine libraries¹⁻³ among them; sponges of the genus *Ircinia* have been proven to be a rich source of diverse biologically active secondary metabolites with novel chemical structures⁴⁻⁷. Several molecules isolated from *Ircinia* species exhibited antineoplastic, antifouling, anti-inflammatory, cytotoxicity⁸⁻¹⁰ and antimicrobial properties¹¹.

Materials and Methods

Experimental details

Optical rotations were determined on a Rudolph Research Analytical (AUTOPOL V) polarimeter at a wavelength of 589 nm (sodium D line) using a 1.0 dm cell with a total volume of 1.0 mL. The UV spectra were measured on an Agilent technologies carry series UV-VIS spectrophotometer and Infrared spectra on Bruker ALPHA. All solvents were of

analytical grade. Column chromatography was performed on Merck silica gel (120-200 mesh) and Sephadex LH-20 (Sigma-Aldrich Chemie GmbH). Thin layer chromatography was carried out with silica gel GF254 plates, Merck, USA. The ¹H and ¹³C, DEPT-135, COSY, TOCSY, HSQC, HMBC at Bruker 400 MHz (or 100 MHz for ¹³C). The chemical shifts (δ) are reported in ppm and coupling constants (J) in Hz. The positive ion HR-ESI-MS spectra were recorded on a Mass Q-TOF-LC-MS spectrometer (Bruker Daltonics).

Collection of sponge

The sponge *Ircinia fusca* (Carter, 1880) was collected from Bagwatibandhan (N 18°19.092, E 072°57.343) West coast of Maharashtra, India in February 2016. The sponge was identified by Dr. Satish S. Mokashe, Associate Professor, Department of Zoology, Dr. Babasaheb Ambedkar Marathwada University, India. A voucher specimen (No. 55-25P) was deposited in the College of Fisheries, Maharashtra, India.

Extraction and isolation

In the laboratory, the sponge was washed with distilled water to remove surface salts, sand, and epiphytes. The sponge was dabbed with tissue paper to remove excess water, cut into small pieces and placed in a lyophilizer to completely dry. The dried material was (2 g dried weight) reduced to small pieces and extracted with MeOH (0.86 g). Desalting of sponge methanolic extract was done with acetone. The methanolic extract was concentrated under vacuum using a rotary evaporator at 40 °C followed by partition between hexane, DCM, water. All the partition layers were subjected to preliminary bioactivity studies (antibacterial and antifungal) by disc diffusion method.

Antimicrobial activity

The isolated compounds 1-3 were tested against antibacterial i.e., *Escherichia coli* (NCIM 2065), *Salmonella typhimurium* (NCIM 2501), *Bacillus subtilis* (NCIM 2063), *Staphylococcus aureus* (NCIM 2079), *Mycobacterium smegmatis* (NCIM 5138) and antifungal strains *Aspergillus niger* (NCIM 1207), *Penicillium chrysogenum* (NCIM 1315), *Alternaria*

*Correspondent author

Email: srinu.meesala@iiserpune.ac.in

Supplementary data available in the online version only.

sp. (NCIM 900), and *Fusarium sp.* (NCIM 1372). The crude extracts were dissolved in DMSO at a concentration of 1 mg/mL. The discs were loaded with different concentrations (10- 500 µg/disk) of the pure compound, to find out the inhibitory potential. The diameters of the inhibition zones generated around the discs were measured (\emptyset in mm). Ampicillin was taken as positive control. The tests were performed in triplicates. DMSO used to dissolve the extracts and the compounds were checked for the absence of antimicrobial activity. The diameters of the halos of inhibition can be rationalized on a qualitative basis as follows: $\emptyset < 7$ mm: inactive, $7 \text{ mm} \leq \emptyset < 8$ mm: slightly active, $8 \text{ mm} \leq \emptyset < 9$ mm: significantly active, $\emptyset \geq 9$ mm: very active. The compound which showed ≥ 9 mm was selected for MIC studies.

Minimum inhibitory concentration (MIC)

Purified compound was evaluated for MIC values against various Gram-positive and Gram-negative bacteria test cultures using nutrient broth described by Andrews¹².

Results and Discussion

During the course of our search for bioactive substances from marine sponges, we collected *Iricnia fusca* from Bagwatibandhan (N 18° 19.092, E 072°57.343), Arabian Sea, West coast of Maharashtra, India. The crude organic methanolic extract of the specimen exhibited antimicrobial activity in preliminary studies. Chromatographic separation of the MeOH extract using C18 semi-preparative reverse phase HPLC led to the isolation of new imidazole derivative (1) as showed in Fig. S1 and two known compounds variabilin (2) and iricinialactam A (3)⁷ (Fig. 1). Herein, the structure elucidation and bioactivity of the new compound 1 from the *I. fusca* is described.

Compound 1 was obtained as a white crystalline solid, melting point 110 °C and $[\alpha]_D^{25} = 13.0$ (c 1.0, CH₃OH). The ESI-MS-QTOF exhibited a pseudo




Fig. 1 — Chemical structure of isolated compounds from *I. fusca*

molecular ion peak at m/z 425.2155 $[M+K]^+$ (Fig. S2), corresponding to the molecular formula of C₁₇H₃₀N₄O₆K, indicating four degrees of unsaturation and four nitrogen atoms in the molecule. The UV absorption at λ_{max} 214 nm indicating the absence of chromophore in the compound 1. The IR spectra showed bands at 3339, 2946, 2034, 1450, 1120, 1027 suggested the presence of hydroxyl, lactam, and aromatic moieties. Its ¹H NMR chemical shifts at δ_H 8.85 (s, H-2), δ_H 7.90 (s, H-5) and ¹³C NMR signals at δ_C 140.0 (C-2), 131.8 (C), and 128.4 (C-5) were a characteristic feature of imidazole ring with C-4 substitution¹³. The ¹H NMR spectrum of compound 1 (Fig.S3) showed three singlet's at δ_H 3.05, 3.91, 4.09 indicating three N-CH₃ groups and one singlet at δ_H 3.41 (6H, s) for two methoxy groups and five methine signals at 3.73 (1H, m, H-2'), δ_H 3.27 (1H, d, $J = 10$ Hz, H-3'), 4.67 (1H, d, $J = 3.7$ Hz, H-3''), 3.60 (1H, d, $J = 10$ Hz, H-5''), 3.38 (1H, dd, $J = 3.7, 10$ Hz, H-4''), and three methylene at δ_H 3.05 (2H, d, $J = 10$ Hz, H-4'), 3.60 (2H, d, $J = 10$ Hz, H-1'), 3.53 (2H, m, H-5'). The ¹³C, DEPT 135, ¹³C HSQC NMR spectrum of 1 (Fig. S4, S5, S6) exhibited total of 17 carbons (Table 1) including a carbonyl signal at δ_C 159.0 (C-2''), four olefinic methines at δ_C 140.0 (C-2), 128.4(C-5), 75.1 (C-5''), 100.8 (C-3''), one quaternary carbon δ_C 131.4(C-4), two methylenes δ_C 50.4 (C-5'), 39.8(C-4'), three oxygenated methines at δ_C 71.8

Table 1 — NMR data of compounds 1 (400 MHz, CD₃OD)

Position	Compound 1	
	δ_H /ppm, multi(J/Hz)	δ_C /ppm
1	N	-
2	8.85, s	140.0, CH
3	N	-
4	-	131.4, C
5	7.90, s	128.4, CH
1'	3.60, d, $J = 10$ Hz	62.7, CH ₂
2'	3.73, m	73.5, CH
3'	3.27, br d, $J = 10$ Hz	71.8, CH
4'	3.05, d, $J = 10$ Hz	39.8, CH ₂
5'	3.53, m	50.4, CH ₂
1''	N	-
2''	-	159.0, CO
3''	4.67, d, $J = 3.7$ Hz	100.8, CH
4''	3.38, brd, $J = 10, 3.7$ Hz	73.5, CH
5''	3.60, d, $J = 10$ Hz	75.1, CH
1''N	4.09, s	36.6
6''N	3.91, s	36.5
1''N	3.05, s	38.4
-	3.41, s	55.5, OCH ₃
-	3.41, s	55.5, OCH ₃

(C-3'), 73.5 (C-2'), 73.5 (C-4''), an oxygenated methylene at δ_C 62.7 (C-1'), two methoxy carbons at δ_C 55.5, three N-methyl carbons at δ_C 36.5, 36.6, 38.4 respectively.

The COSY (Fig.S7) correlations between H-4''/H-3'', H-4''/H-5', as well as the HMBC (Fig.S8) correlations from H-3'' to C-2'' revealed the presence of pyrrolidin-2-one ring moiety. Further HMBC correlations of N-methyl proton δ_H 3.05 to C-2'', δ_H 3.41 (Methoxy) with C-3'', C-5'' asserted the location of the methoxy groups at H-3'' and H-5'' positions respectively as shown in Fig. 2, which confirmed 3, 5 dimethoxy -1-methyl pyrrolidin-2-one moiety in compound 1. The 1H signals at δ_H 3.54 and 3.68 and correlations of H-5'-H-4'-H-3'-H-2'-H-1' observed in the COSY spectrum, in combination with HMBC correlations of H-5' to N-methyl carbon δ_C 36.6 (C-6') revealed the presence of amino pentyl moiety in compound 1. Furthermore, HMBC correlations of 6'N to C-4 and H-3' to C-4'' revealed 3, 5 dimethoxy -1-methyl pyrrolidin-2-one was linked to 1-methyl-1H-imidazol-4-yl-amino-pentan-3-yl-oxy through ether linkage showed in Fig. 2. Thus, the structure compound **1** was established as 4-((1, 2-dihydroxy-5-(methyl (1-methyl-1H-imidazol-4-yl) amino) pentan-3-yl) oxy)-3,5-dimethoxy-1-methylpyrrolidin-2-one as shown in Fig. 1

Compound 1 to 3 were evaluated for their antifungal activity against *A. niger* (NCIM 1207), *P. chrysogenum* (NCIM 1315), *Alternaria sp* (NCIM 900), *Fusarium sp* (NCIM 1372) as well as antibacterial activity against *E. coli* (NCIM 2065), *S. typhimurium* (NCIM 2501), *B. subtilis* (NCIM 2063) and *S. aureus* (NCIM 2079) was also tested using the disc diffusion assay¹⁴. Neither antifungal, nor antibacterial activity was recorded for the compounds (2 and 3), but compound 1 exhibited selective inhibitory growth activity against *S. aureus* when compared with ampicillin disc at 100 $\mu\text{g/mL}$. The MIC of compound 1 was found to be 280 μM . This indicates its use as a potential drug against gram-positive bacteria which causes nosocomial infections in hospitals, especially *S. aureus*. We are pursuing the activity of this drug against methicillin-resistant *Staphylococcus aureus* strains.




Fig. 2 — Key COSY and HMBC correlations of compound 1

Conclusion

Until now, sponge species like *Leucetta*¹⁵ and *Leucosolenia*¹⁶ have been reported as rich sources of imidazole alkaloids. This is the first report that the sponge *Iricinia fusca*, taken from the West coast of Maharashtra, produces a new imidazole alkaloid as a natural product. The isolated compound **1** was similar to 2,5-Pyrrolidinedione, 1,1'-(1,4-butanediyl)bis[3-ethoxy-) with CAS Registry Number 1403656-78-9, which was found in SciFinder to be synthesized via a chemical reaction.

In summary, three compounds were isolated from *I. fusca*. Compound **1** has been reported as a new alkaloid metabolite isolated from *I. fusca* and the structure was elucidated by NMR and mass spectroscopic analysis. The genus *Iricinia* has been known to produce diverse metabolites like sesterterpenoids, irciniastatins, quinones, ircinialactams, and pyrroles, but the current study also showed the presence of new imidazole derivative from *I.fusca*. These reports indicate that a large resource of metabolites with biological significance may yet be disclosed in *I. fusca*.

Compound 1: 1H NMR (400 MHz, MeOD) δ 8.85 (s, 1H), 7.90 (s, 1H), 4.67 (d, $J = 3.7$ Hz, 1H), 4.09 (s, 3H), 3.91 (s, 3H), 3.73 (m, 1H), 3.60 (d, $J = 10$ Hz, 3H), 3.53 (m, 2H), 3.38 (br d, $J = 10, 3.7$ Hz, 1H), 3.41 (s, 6H), 3.27 (d, $J = 10$ Hz, 1H), 3.05 (d, $J = 10.0$ Hz, 5H).

Acknowledgement

This work was financially supported (RGSTC/File-2007/DPP-054/CR-28) by the Rajiv Gandhi Science and Technology Commission, Maharashtra, India. We are thankful to NMR Research Centre and Mass facility at IISER Pune. The authors thank NRC, IISc, Bangalore for recording NMR experiments and Sandeep C. Kanade for mass services.

Conflict of Interest

The authors declare no competing financial interests.

References

- Faulkner D J, Marine natural products, *Nat Prod Rep*, 2002, **19**, 1-49.
- Blunt J W, Copp B R, Ping Hu W, Munro M H G, Northcote P T, *et al.*, Marine natural products, *Nat Prod Rep*, 2009, **26**, 170-244.
- Blunt J W, Copp B R, Keyzers R A, Munro M H G and Prinsep M R, Marine natural products, *Nat Prod Rep*, 2015, **32**, 116-211.
- Malcolm S B, Annette E, Gordon K, Jacqueline W and Ronald J Q, Cheilanthane sesterterpenes, protein kinase inhibitors, from a marine Sponge of the genus *Iricinia*, *J Nat Prod*, 2001, **64**, 300-303.

- 5 George R P, Jun-Ping X, Jean-Charles C, Robin K P, Larry P T, *et al.*, Antineoplastic agents. 520. Isolation and structure of Irciniastatins A and B from the Indo-pacific marine sponge *Ircinia ramosa*, *J Med Chem*, 2004, **47**, 1149–1152.
- 6 Maria T, Claire H, Constantinos V, Catherine H and Vassilios R, Chemical defense and antifouling activity of three Mediterranean sponges of the genus *Ircinia*, *Z Naturforsch*, 2002, **57c**, 161-171.
- 7 Walter B, Robiul I, Frank F, Andrew M P, Hua Z, *et al.*, Ircinialactams: Subunit-selective glycine receptor modulators from Australian sponges of the family Irciniidae, *Bioorganic Med Chem*, 2010, **18**, 2912–2919.
- 8 Rashid M A, Gustafson K R and Boyd M R, New chondropsin macrolide lactams from marine sponges in the genus *Ircinia*, *Tetrahedron Lett*, 2001, **42**, 1623-1626.
- 9 Yan S, Zhang G, Su J and Zeng L, A rare long conjugated diterpene ketene from the marine sponge *Ircinia selaginea* (Lamarck), *Gaodeng Xuexiao Huaxue Xuebao*, 2001, **22**, 949-951.
- 10 Kokubo S, Yogi K, Udin M J, Inuzuka T, Suenaga K, *et al.*, Kohamaic acids A and B, novel cytotoxic sesterterpenic acids, from the marine sponge *Ircinia sp.*, *Chem Lett*, 2001, **30**, 176-177.
- 11 Sharad S, Satish M, Gupta S G, Vijay L and Dinesh K, Antimicrobial potential of the marine intertidal sponge, *Ircinia fusca* from West coast of India, *Indian J Pharm Biol Res*, 2015, **3**, 12-14.
- 12 Andrews J M, Determination of minimum inhibitory concentrations, *J Antimicrob Chemother*, 2001, **48**, 5-16.
- 13 Shin J, Rho J R, Seo Y, Lee H S, Cho K W, *et al.*, Wondonins A and B, new bis(dihydroxystyryl)imidazoles from a two-sponge association, *Tetrahedron Lett*, 2001, **42**, 1965–1968.
- 14 Lippert H, Brinkmeyer R, Mulhaupt T and Iken K, Antimicrobial activity in sub-Arctic marine invertebrates, *Polar Biol*, 2003, **26**, 591-600.
- 15 Tsukamoto S, Kawabata T, Kato H, Ohta T, Rotinsulu H, *et al.*, Naamidines H, Cytotoxic Imidazole alkaloids from the Indonesian Marine sponge *Leucetta chagosensis*, *J Nat Prod*, 2007, **70**, 1658–1660
- 16 Ralifo P, Tenney K, Valeriote F A and Crews P, A distinctive structural twist in the aminoimidazole alkaloids from a calcareous Marine sponge: Isolation and characterization of Leucosolenamines A and B, *J Nat Prod*, 2007, **70**, 33-38.